

ORIGINAL ARTICLE

# Influence of Nitrogen Fertilizer and Vermicompost Application on Flower Yield and Essential Oil of Chamomile (*Matricaria Chamomile* L.)

Mohammad Reza Haj Seyed Hadi\*, Mohsen Abarghoeei Fallah, Mohammad Taghi Darzi

Department of Agronomy, Islamic Azad University, Roudehen Branch, Roudehen, Iran

(Received: 30 April 2015 Accepted: 1 July 2015)

## KEYWORDS

Vermicompost;  
Chamomile;  
Nitrogen;  
Essential oil

**ABSTRACT:** This study was performed to assess the effects of nitrogen fertilizer and vermicompost on qualitative and quantitative yield of chamomile (*Matricaria chamomilla* L.). It was conducted at the Research Fields of Ran Company located in Firouzkouh, Iran, in 2013. Treatments were consisted of 1) Control, 2) 100% nitrogen from urea, 3) 100% nitrogen from ammonium nitrate, 4) 75% nitrogen from urea and 25% from vermicompost, 5) 75% nitrogen from ammonium nitrate and 25% from vermicompost, 6) 50% nitrogen from urea and 25% from vermicompost, 7) 50% nitrogen from ammonium nitrate and 25% from vermicompost, 8) 25% nitrogen from urea and 25% from vermicompost, 9) 25% nitrogen from ammonium nitrate and 25% from vermicompost, and 10) 100% nitrogen from vermicompost. The maximum plant height (67.03 cm) and plant weight (93.21 g/plant) were obtained at N2 treatment (200 kg ha<sup>-1</sup> urea). N5 treatment (202.5 kg ha<sup>-1</sup> ammonium nitrate + 1.5 ton vermicompost ha<sup>-1</sup>) caused maximum flower diameter. The highest fresh flower yield (7539.45 kg ha<sup>-1</sup>), dry flower yield (1715.93 kg ha<sup>-1</sup>) and essential oil yield (6.95 kg ha<sup>-1</sup>) obtained in plots, which received 135 kg ha<sup>-1</sup> nitrate ammonium + 3 ton vermicompost ha<sup>-1</sup>. It seems using biofertilizers such as vermicompost could enhance quantitative and qualitative characteristics of chamomile. Moreover, by substituting chemical fertilizers by biofertilizers, ecosystem health and quality of life will increase which it is the most important goals of sustainable developments.

## INTRODUCTION

Chamomile (*Matricaria chamomilla* L.) is a plant from Asteraceae family [1, 2]. Plant height ranges from 20 to

50 cm, according to the ecological conditions of the production region. Root system is short and widespread

\* Corresponding author: mrhshadi@yahoo.com (M.R. Haj Seysd Hadi).

mainly in top 30 cm layer of soil [3]. Because of medicinal properties, its production and consumption is of high interest in most countries [4]. The growth indices, flower yield and active substance of chamomile plants are affected by genetic background and environmental conditions. These attributes are strongly modified by many other factors and management skills such as using the different kinds of fertilizers [1, 5].

In recent decades, production of agricultural crops has been mainly reliant on use of chemical inputs, resulted in environmental problems. One of the approaches to alleviate this problem is to use the strategies, which are based on the principles of organic farming in agro-systems. In such system, synthetic inputs e.g. chemical fertilizers and pesticides are replaced with the use of crop rotation with pulse crops, plant debris, and different types of manures, organic and biological fertilizers. This results in control of weeds and pests and increase of biodiversity in the fields while conserving nutrients in the soil [6].

Growth and qualitative characteristics of chamomile depend on management of nutrients [7, 8]. Among the macro-elements, nitrogen has the most significant effect on growth characteristics of chamomile [9, 10]. Increasing concerns about the negative effects of chemical fertilizers have generated substantial interest in the use of other N sources [11- 13]. Despite the importance of nitrogen in production of field crops and medicinal plants, application of this nutrient element from chemical sources has resulted in environmental problems and pollution of the global ecosystem [14]. One of the strategies to alleviate this problem is to use the methods, based on the principles of organic farming in agro-systems [6].

Sustainable agriculture aims at considerable reduction in application of chemical inputs by following the ecological principles and so it is regarded a proper solution to overcome the current agricultural problems [15].

Conventional agriculture with focus on chemical fertilizers has caused rapid reduction in organic matter and fertility of soils. Therefore, it is essential to apply proper production managements in order to improve physical, chemical and biological characteristics of soils [16].

Application of biofertilizers is among the approaches of crop fertilization to achieve the objectives of sustainable agriculture [17, 18]. Biofertilizers are the bacterial and fungal microorganisms and their metabolic compounds. Biofertilizers contain the conserving materials of high density derived from one or many helpful soil-borne microorganisms or their metabolites that form clones in root peripheral or plant internal sections and induce the host plant growth by numerous strategies [19, 20].

One of the most important biological fertilizers produced by earthworms (especially *Eisenia foetida*), is vermicompost. Vermicompost is rich in macro- and microelements that are in available forms for plants [21]. High microbial population and their diversity, containing plant growth regulators or hormones and large amounts of humic acids, are the advantages of vermicompost [21-23].

Vermicompost could increase plant weight [24] and nutrient use efficiency [25]. There are many reports on positive effects of vermicompost on field crops [26- 29]. Growth, yield and active substance of some medicinal plants such as basil [30], garlic [31], fennel [32], chamomile [2, 33] and roman chamomile [34] were increased by vermicompost application.

This study was done to assess the effects of nitrogen fertilizer and vermicompost on qualitative and quantitative yield of chamomile (*M. chamomilla* L.).

## MATERIALS AND METHODS

### *Field experiment*

The investigation was conducted at the Research Fields of Ran Company located in Firouzkouh (Latitude: 35° 45' N; Longitude: 52° 45' E; Altitude: 1930 m), Iran, in

spring of 2013. Before seed sowing, soil of the experiment site was sampled randomly at top 0-30 cm layer of the field in crinkle method and analyzed at Khak

Azmoon Pishahang Laboratory located at Ghom Province (Table 1).

**Table 1.** Results of soil analysis for the experimental field.

Texture	Clay	Silt	Sand	ECe(dS/m)	pH	OC%	N%	P(ppm)	K(ppm)
Loamy-Clay	34	24	42	3.39	7.6	0.65	0.055	10	300

The statistical design was a randomized complete block design (RCBD) with three replications. Treatments were consisted of:

- 1- Control
- 2- 100% nitrogen from urea (200 kg ha<sup>-1</sup>)
- 3- 100% nitrogen from ammonium nitrate (270 kg ha<sup>-1</sup>)
- 4- 75% nitrogen from urea and 25% from vermicompost (150 kg ha<sup>-1</sup> urea + 1.5 ton vermicompost ha<sup>-1</sup>)
- 5- 75% nitrogen from ammonium nitrate and 25% from vermicompost (202.5 kg ha<sup>-1</sup> ammonium nitrate + 1.5 ton vermicompost ha<sup>-1</sup>)
- 6- 50% nitrogen from urea and 25% from vermicompost (100 kg ha<sup>-1</sup> urea + 3 ton vermicompost ha<sup>-1</sup>)

- 7- 50% nitrogen from ammonium nitrate and 25% from vermicompost (135 kg ha<sup>-1</sup> ammonium nitrate + 3 ton vermicompost ha<sup>-1</sup>)
- 8- 25% nitrogen from urea and 25% from vermicompost (50 kg ha<sup>-1</sup> urea+4.5 ton vermicompost ha<sup>-1</sup>)
- 9- 25% nitrogen from ammonium nitrate and 25% from vermicompost (67.5 kg ha<sup>-1</sup> ammonium nitrate + 4.5 ton vermicompost ha<sup>-1</sup>)
- 10- 100% nitrogen from vermicompost (6 ton vermicompost ha<sup>-1</sup>)

The nitrogen requirements of chamomile is 90 kg N /ha [35]. According to this value, treatments were calculated. Chemical analysis of vermicompost is presented in Table 2.

**Table 2.** Results of vermicompost analysis.

pH	ECe ds/m	N%	K%	P%	OC%	OM%	Moisture%
7	1.1	4.92	3.19	0.61	37.7	65	25

Vermicompost was applied to the top 5-10 cm layer of soil. Experimental plots were 3 × 2.25m. Seeds were obtained from Medicinal Plants Research Center, Isfahan, Iran. Seeds were sown on the rows with 30 cm spacing. 15-20 days after sowing, the plants were thinned to 10 cm distance between plants. Irrigated was done weekly.

### Measurements

Measured traits included plant height (cm), flower head diameter (mm), plant weight (g plant<sup>-1</sup>), flower yield (kg

ha<sup>-1</sup>) and essential oil percentage. Samplings were done on the middle rows in each plot.

Plant height was measured by a ruler at the flowering stage. Flower head diameter was measured using a vernier caliper. 1m<sup>2</sup> of inner rows was harvested for calculating flower yield. Dry weight of flowers was measured after putting the plants at room temperature (25° C for 72 h) [36].

Essential oil extraction was carried out at the IBB Laboratory, University of Tehran, using a Clevenger-type apparatus. Protocol of Letchamo and Marquard [29] was applied for measuring essential oil in the flowers.

To determine the amount of essential oil, a sample of 100 g of flowers was mixed with 500 ml of tap water in a flask and the water was distilled for 3 h using a Clevenger-type apparatus. The oil content was measured by following the protocol of Letchamo and Marquard [37], based on ml oil per 100 g dry matter of flower. Essential oil yield was obtained by multiplying essential oil percentage and dry flower yield.

SAS 9.1 software was used to analyze the data [38]. Means comparisons were carried out by Duncan's Multiple Range Test at 5% probability level.

## RESULTS

The results of analysis of variance showed that all the measured traits were significantly affected by the vermicompost and nitrogen fertilizer (Table 3).

**Table 3.** Analysis of variance for the measured traits of chamomile (*Matricaria chamomilla* L.).

S.O.V	df	Plant Height	Plant Weight	Flower Head Diameter	Fresh Flower Yield	Dry Flower Yield	Essential Oil Content	Essential Oil Yield
Rep	2	148.800333**	1610.006470**	0.490033	2130755.40	99558.299	0.00089333	0.20019000
Treat	9	214.261815**	819.261815**	14.5832741*	6345953.11*	257575.206**	0.02098667**	7.47748296**
Error	18	15.994037	49.08403	5.7196219	1837899.01	59764.971	0.00047111	0.80997519
CV%		7.47	10.14	13.09	14.80	19.93	6.41	20.41

### *Plant height*

Chamomile plant height was significantly affected by various levels of treatments. The maximum height (67.03 cm) was obtained by using 200 kg ha<sup>-1</sup> urea (Table 4).

### *Flower head diameter*

Flower head diameter was significantly affected by treatments. Use of 198.75 kg ha<sup>-1</sup> ammonium nitrate + 1.5 ton vermicompost ha<sup>-1</sup> caused the maximum flower head diameter (Table 4).

### *Pant weight*

Fresh and dry weights of chamomile plants were significantly affected by treatments. The maximum fresh weight (93.207 g plant<sup>-1</sup>) and dry weight (11.843 g plant<sup>-1</sup>) were obtained when 200 kg ha<sup>-1</sup> urea was used (Table 4).

### *Fresh and dry flower yield*

Plants received ammonium nitrate and vermicompost simultaneously had significantly greater flower yield. Flower yields (7539.2 and 1715.9 kg ha<sup>-1</sup> for fresh and dry yields, respectively) were maximum at 132.5 kg ha<sup>-1</sup> ammonium nitrate + 3 ton vermicompost ha<sup>-1</sup> (Table 4; Figure 1).




Figure 1. Effects of treatments on fresh flower yield.

### Essential oil

Essential oil content was influenced by urea, ammonium nitrate and vermicompost application. The maximum

essential oil yield ( $6.95 \text{ kg ha}^{-1}$ ) (Fig. 2) was obtained by using  $6 \text{ ton ha}^{-1}$  vermicompost.




Figure 2. Effects of treatments on essential oil content in chamomile flowers.

**Table 4.** Mean comparison for treatment effects on measured traits for chamomile.

Treatments	Plant Height (cm)	Plant Weight (g/Plant)	Flower Head Diameter (mm)	Fresh Flower Yield (kg/ha)	Dry Flower Yield (kg/ha)	Essential Oil Content (%)	Essential Oil Yield (kg/ha)
1	37.66 e	35.46 f	13.19 b	2346.11 c	661.60 d	0.21 f	1.38 e
2	67.03 a	93.21 a	18.52 a	5489.21 ab	1296.21 abc	0.33 c	4.23 cd
3	60.47 ab	77.14 bc	17.49 ab	5352.10 ab	1291.32 abc	0.30 cd	3.89 d
4	59.90 ab	70.69 bcd	16.78 ab	5564.72 ab	1320.11 abc	0.26 e	3.41 d
5	58.96 bc	70.38 cd	20.90 a	4706.84 b	1134.48 c	0.31 cd	3.49 d
6	58.73 bc	52.03 bc	20.19 a	6393.20 ab	1450.10 abc	0.28 de	4.05 cd
7	52.37 cd	83.65 ab	18.53 a	7539.45 a	1715.93 a	0.41 b	6.95 a
8	46.43 d	52.02 e	19.89 a	7243.96 ab	1614.18 ab	0.38 b	6.06 ab
9	47.16 d	66.87 cd	17.60 ab	5191.44 ab	1098.30 c	0.45 a	4.96 bcd
10	46.30 d	61.58 de	19.45 a	4812.27 b	1221.77 bc	0.46 a	5.64 bc

1- 1) Control, 2) 100% nitrogen from urea (200 kg ha<sup>-1</sup>), 3) 100% nitrogen from nitrate ammonium (265 kg ha<sup>-1</sup>), 4) 75% nitrogen from urea and 25% from vermicompost (150 kg ha<sup>-1</sup> urea + 1.5 ton vermicompost ha<sup>-1</sup>), 5) 75% nitrogen from nitrate ammonium and 25% from vermicompost (198.75 kg ha<sup>-1</sup> nitrate ammonium + 1.5 ton vermicompost ha<sup>-1</sup>), 6) 50% nitrogen from urea and 25% from vermicompost (100 kg ha<sup>-1</sup> urea + 3 ton vermicompost ha<sup>-1</sup>), 7) 50% nitrogen from nitrate ammonium and 25% from vermicompost (132.5 kg ha<sup>-1</sup> nitrate ammonium + 3 ton vermicompost ha<sup>-1</sup>), 8) 25% nitrogen from urea and 25% from vermicompost (50 kg ha<sup>-1</sup> urea + 4.5 ton vermicompost ha<sup>-1</sup>), 9) 25% nitrogen from nitrate ammonium and 25% from vermicompost (66.25 kg ha<sup>-1</sup> nitrate ammonium + 4.5 ton vermicompost ha<sup>-1</sup>), 10) 100% nitrogen from vermicompost (6 ton vermicompost ha<sup>-1</sup>)

## DISCUSSION

According to the results, 200 kg ha<sup>-1</sup> urea increased the plant height and plant weight. Organic amendments only release 30-35% of total nitrogen in the first growing season. It seems nitrogen uptake by plants from chemical sources is more than that from biofertilizers such as vermicompost [35]. It was found that nitrogen enhances vegetative growth and plant height of chamomile [39].

Nitrogen application had significant effect on the number of flowers per plant, number of auxiliary branches, and flower yield [40]. Nitrogen had a significant effect on plant height, number of flowers per plant, and dry flower yield of chamomile [41]. The positive effect of nitrogen on root characteristics and essential oil content and chamazulene of chamomile has also been reported [14, 42 and 44].

Application of biofertilizers such as vermicompost in a sustainable agriculture system improves the yield and quality of active ingredient in medicinal plants [15, 17, 45, and 46]. The impact of different levels of vermicompost and irrigation were investigated on

morphological traits and essential oil content of chamomile and found that increased level of vermicompost significantly improved the plant height, early flowering, flower yield, length, and diameter of receptacle [36]. Results of another research conducted on basil (*Ocimum basilicum* L.) revealed that application of vermicompost improved the yield and quality of essential oil content, essential oil yield and biological yield as compared with the control treatment [30]. Furthermore, the results of research conducted on one of the species of *Davana* (*Artemisia pallens*) indicated that application of vermicompost significantly improved the essential oil yield as compared with the control treatment and this was due to the increase of dry matter content [11]. Arguello et al. [47] displayed the improved quality of garlic (*Allium sativum*) because of vermicompost application. The increase of yield and quality of rose-scented geranium (*Pelargonium graveolens*) were shown due to the application of vermicompost [48]. Application of 10 ton vermicompost increased the essential oil contents in Fennel

(*Foeniculum vulgare*) [49]. Similar results have been reported on the positive effects of vermicompost on the medicinal plant garlic [31], coriander [32], chamomile [33] and Roman chamomile [34].

Using vermicompost and chemical nitrogen together resulted in maximum values, which are in accordance with Fallahi et al. [50]. Vermicompost could increase plant growth and development by improving physical, chemical and microbial properties of soil [50]. Vermicompost is a main factor for increasing flower yield of chamomile and marigold [51]. Integration of chemical nitrogen and vermicompost increases plant growth indices such as leaf area index, roots length and plant dry weight [52].

Essential oil is a terpenoid compound and its components (isoprenoids) such as Isopantyl pyrophosphate (IPP) and Dimethyl Alyl pyrophosphate (DMAPP) highly demand NADPH, ATP, and the presence of elements such as nitrogen and phosphorous is necessary for formation of the latter compounds. Therefore, while interpreting the results of improved essential oil content caused by application of vermicompost, it could be stated that the increased vermicompost can increase the essential oil content of flower by increasing the absorption of nitrogen and phosphorous [35].

It was found that applying 5 ton vermicompost along with 50, 25 and 25 kg ha<sup>-1</sup> of N, P and K, respectively, increase essential oil percentage, essential oil yield and biological yield of basil [30]. Other investigations have shown that vermicompost increases the essential oil of chamomile [33, 53].

### CONCLUSIONS

The maximum plant height and plant weight were observed for using 200 kg ha<sup>-1</sup>. Flower yield and essential oil yield of chamomile plants cultivated in soil amended with urea were minimum. The maximum values for these traits were recorded at plots, which

received 135 kg ha<sup>-1</sup> ammonium nitrate + 3 ton vermicompost ha<sup>-1</sup>. However, the maximum essential oil content obtained by applying 6 ton vermicompost ha<sup>-1</sup>. The application of the vermicompost had positive effect on essential oil and by increasing vermicompost rate essential oil of flowers increased.

In the case of chamomile (and of course all medicinal plants) the quality and thus the concentration of essential oil and its yield are much more relevant than the flower yield. As our results showed, essential oil content was maximum by applying 6 ton vermicompost ha<sup>-1</sup>, while the maximum essential oil yield was related to N7 treatment (135 kg ha<sup>-1</sup> ammonium nitrate + 3 ton vermicompost ha<sup>-1</sup>).

Integration of vermicompost and ammonium nitrate fertilizer could increase growth, flower yield and essential oil of chamomile and have thus important roles for sustainable production of chamomile.

### ACKNOWLEDGMENTS

The authors wish to thank Eng. Mohammad Reza Bahrehvar, Ran Agricultural Company Manager for his technical cooperation during this study. The authors declare that there is no conflict of interests.

### REFERENCES

1. Mosleh Z., Salehi M.H., Rafiei Alhosseini M., 2014. Effects of chamomile cultivation and manure on silt and clay changes in soils. *J Agric Environ.* 18(68), 11-21.
2. Dastgheibifard N., Sharafzadeh S., Bazrafshan F., 2014. Influence of cow manure vermicompost on growth characteristics of German chamomile. *Cibtech J Zool.* 3(1), 58-61.
3. Sharafzadeh S., Alizadeh O., 2011. German and Roman chamomile. *J App Pharm Sci.* 1(10), 1-5.
4. Baghalian K., 2000. Recent advances of medicinal and aromatic plants in Iran. *Chronica Hort.* 40, 13–14.

5. Ghani A., Azizi M., Hasanzadeh Khayyat M., Pahlavanpour A., 2008. Essential oil composition of *Achillea eriophora*, *A. nobilis*, *A. biebersteinii* and *A. wilhelmsii* from Iran. *J Essent Oil Bear Pl.* 11(5), 460-467.
6. Elsen T.V., 2000. Species diversity as a task for organic agriculture in Europe. *Agr Ecosyst Environ.* 77, 101-109.
7. Emonger V.E., Chweya J.A., 1992. Effect of nitrogen and variety on essential oil yield and composition of chamomile flowers. *Trop Agr.* 69, 290-292.
8. Heneka N., 1993. *Chamomilla recutita* L. *Aust J Med Herbal.* 5, 33-39.
9. Letchamo W., 1992. A comparative study of chamomile yield, essential oil and flavonoids content under two sowing seasons and nitrogen levels. *Acta Hort.* 306, 375-384.
10. Karami A., Khush-Khui M., Sefidkon F., 2008. Effects of nitrogen, phosphorous and potassium on yield, essential oil content and composition of cultivated and wild population of *Chamomilla recutita* L. *Rauschert. J Med Aroma Plant Sci.* 30, 1-4.
11. Pandey R., 2005. Mangement of *Meloidogyne incognita* in *Artemisia pallens* with bio-organics. *Phytoparasitica.* 33(3), 304-308.
12. Ashwini K.M., Sridhar K.R., 2006. Evaluation of pill millipede (*Arthrosphaera magna*) compost on plant growth and dry matter yield. *J Environ. Agr Food Chem.* 5, 1323–1329.
13. Najm A.A., Haj Seyed Hadi M.R., Fazeli F., Darzi M.T., Rahi A., 2012. Effect of integrated management of nitrogen fertilizer and cattle manure on the leaf chlorophyll, yield, and tuber glycoalkaloids of *Agria* potato. *Commun Soil Sci Plant.* 43(6), 912-923.
14. Nikolova A., Kozhuharova K., Zheljazkov V.D., Craker L.E., 1999. Mineral nutrition of chamomile (*Chamomilla recutita* L.). *Acta Hort.* 502, 203-208.
15. Darzi M.T., Haj Sayed Hadi M.R., Rejali F., 2012. Effects of the application of vermicompost and nitrogen fixing bacteria on quantity and quality of the essential oil in dill (*Anethum graveolens*). *J Med Plants Res.* 6(21), 3793-3799.
16. Liu M., Hu F., Chen X., Huang Q., Jiao J., Zhang B., Li H., 2009. Organic amendments with reduced chemical fertilizer promote soil microbial development and nutrient availability in a subtropical paddy field: the influence of quantity, type and application time of organic amendments. *Appl Soil Ecol.* 42, 166-175.
17. Kapoor R., Giri B., Mukerji K.G., 2004. Improved growth and essential oil yield and quality in *Foeniculum vulgare* Mill on mycorrhizal inoculation supplemented with P-fertilizer. *Bioresource Technol.* 93, 307-311.
18. Shaalan M.N., 2005. Effect of compost and different sources of biofertilizers, on borage plants (*Borago officinalis* L.). *Egypt J Agr Res.* 83(1), 271-284.
19. Omidi H., Naghdibadi H.A., Golzad A., Torabi H., Fotookian M.H., 2009. Effects of chemical and biological nitrogen fertilizers on quantitative and qualitative yield of saffron (*Crocus sativus* L.). *J Med Plants.* 9(30), 98-109.
20. Manafee W.F., Kloepper J.W., 1994. Application of plant growth promoting rhizobacteria in sustainable agriculture. p. 23-31. C.E. Paankburst et al. (ed.) *Soil biota management in sustainable farming systems.* CSIRO, Pub. East Melbourne, Australia.
21. Sadeghi A.A., Klarestaghi K., Ghalibaf K., 2014. Effects of urea fertilizer and vermicompost on yield of *Althea officinalis* L. *J Agroecol.* 6(1), 42-50.
22. Heidarianpoor M.B., Sameni A., Sheikhi J., Karimian N., Zareie M., 2014. Effects of vermicompost and nitrogen on vegetative growth and nutrient uptake by sunflower. *J Agr Environ.* 18(67), 217-227.
23. Gutierrez-Miceli F.A., Santiago-Borraz J., Montes Molina J.A., Carlos Nafate C., Abud-Archila M., Oliva Laven M.A., Rincon-Rosales R., Dendooven L., 2007. Vermicompost as a soil supplement to improve growth, yield and fruit quality of tomato (*Lycopersicum esculentum*). *Bioresource Technol.* 98, 2781-2786.



24. Ali M., Griffiths A.J., Williams K.P., Jones D.L., 2007. Evaluating the growth characteristics of lettuce in vermicompost and green waste compost. *J Soil Biol.* 43, S316-S319.
25. Chaoui H.I., Zibilske L.M., Ohno T., 2003. Effects of earthworm casts and compost on soil microbial activity and plant nutrient availability. *Soil Biol Biochem.* 35, 295-302.
26. Singh R., Sharma R.R., Kumar S., Gupta R.K., Patil R.T., 2008. Vermicompost substitution influences growth, physiological disorders, fruit yield and quality of strawberry (*Fragaria x ananassa* Duch.). *Bioresource Technol.* 99, 8507-8511.
27. Fernandez-Luqueno F., Reyes-Valera V., Martinez-Suarez C., Salmon-Hernandez G., Yanez-Menese J., Ceballos-Ramirez J.M., Dendooven L., 2010. Effect of different nitrogen sources on plant characteristics and yield of common bean (*Phaseolus vulgaris* L.). *Bioresource Technol.* 101, 396-403.
28. Suthar S., 2009. Impact of vermicompost and composted farmyard manure on growth and yield of garlic (*Allium stivum* L.) field crop. *Int J Plant Prod.* 3(1), 27-38.
29. Najar I.A., Khan A.B., 2013. Effect of Vermicompost on Growth and Productivity of Tomato (*Lycopersicon esculentum*) Under Field Conditions. *Acta Biol Malaysiana.* 2(1), 12-21.
30. Anwar M., Patra D.D., Chand S., Alpesh K., Naqvi A.A., Khanuja S.P.S., 2005. Effect of organic manures and inorganic fertilizer on growth, herb and oil yield, nutrient accumulation, and oil quality of French basil. *Commun. Soil Sci Plant.* 36(13-14), 1737-1746.
31. Verma S., Choudhary M.R., Yadav B.L., Jakhar M.L., 2013. Influence of vermicompost and sulphur on growth and yield of garlic (*Allium sativum* L.) under semi-arid climate. *J Spice Aroma Crops.* 22(1), 20-23.
32. Godara A.S., Gupta U.S., Lal G., Singh R., 2014. Influence of organic and inorganic source of fertilizers on growth, yield and economics of coriander (*Coriandrum sativum* L.). *Int J Seed Spice.* 4(2), 77-80.
33. Ansarifar M., Noormohamadi G., Seyed Hadi M.R.H., Riazi G., 2012. Effect of organic nutrients on flower yield and oil content of chamomile (*Matricaria chamomilla* L.). *J Med Plants By-products.* 2, 177-181.
34. Sirousmehr A., Arabi J., Asgharipour M.R., 2014. Effect of drought stress levels and organic manures on yield, essential oil content and some morphological characteristics of sweet basil (*Ocimum basilicum*). *Adv Environ Biol.* 8(4), 880-885.
35. Ghazi Manas M., Banj Shafiee S., Haj Seyed Hadi M.R., Darzi M.T., 2013. Effects of vermicompost and nitrogen on quantitative and qualitative yeild of chamomile (*Matricaria chamomilla* L.). *Iran J Med Aroma Plants.* 29(2), 269-280.
36. Azizi M., Rezwane F., Hassanzadeh Khayat M., Lakzian A., Neamati H., 2008. The effect of different levels of vermicompost and irrigation on morphological properties and essential oil content of German chamomile (*Matricaria recutita* variety Goral). *Iran J Med Aroma Plants.* 24(1), 82-93.
37. Letchamo W., Marquard R., 1993. The pattern of active substances accumulation in chamomile genotypes under different growing conditions and harvesting frequencies. *Acta Hort.* 331, 357-364.
38. SAS Inc., 2002. SAS Procedure guide. SAS Institute, Cary, NC. Schank.
39. Niknejad M., Lebaschy M.H., Jaimand K., Hatami F., 2013. Effect of organic and chemical fertilizers on essential oil of *Matricaria chamomilla* L. *Iran J Med Aroma Plants.* 29(2), 373-386.
40. Zeinali H., Moslehi Yazddeli A., Safaee L., Jaberlansar Z., Akhoondi A., Eskandari Z., 2014. Effects of Nitrogen, Phosphorous and Potassium fertilizers on quantitative and qualitative characteristics of chamomile (*Matricaria chamomilla* L.). *Iran J Med Aroma Plants.* 30(4), 511-518.
41. Dadkhah A., Amini Dehaghi M., Kafi M., 2012. Effects of nitrogen and phosphorous fertilizers on quantitative and qualitative yield of chamomile. *Iran J Field Crop Res.* 10(2), 321-326.

42. Rahmati M., Azizi M., Hasanzadeh Khayyat L., Nemati H., 2009. Effects of plant density and nitrogen on morphological traits, yield, essential oil and chamazulene content of chamomile. *J Hort Sci.* 23, 27-35.
43. Zeinali H., Poorjafa M., Bagheri M., Golparvar M.R., Shiranirad A., 2008. Effects of planting time and nitrogen fertilizer on yield and yield components of chamomile. *Iran J Agron.* 3(10), 220-230.
44. Jadwiga Andrzejewska J., Woropaj-Janczak M., 2014. German chamomile performance after stubble catch crops and response to nitrogen fertilization. *Ind Crop Prod.* 62: 350-358.
45. Ratti N., Kumar S., Verma H.N., Gautam, S.P., 2001. Improvement in bioavailability of tricalcium phosphate to *Cymbopogon martini* var. *motia* by rhizobacteria, AMF and *Azospirillum* inoculation. *Microb Res.* 156, 145-149.
46. Sharma, A.K., 2002. Biofertilizers for sustainable agriculture. *Agrobios*, India, 407 p.
47. Arguello J.A., Ledesma A., Nunez S.B., Rodriguez C.H., Goldfarb, M.D.D., 2006. Vermicompost effects on bulbing dynamics, nonstructural carbohydrate content, yield, and quality of Rosado paraguayo garlic bulbs. *Hort Sci.* 41(3), 589-592.
48. Chand S., Pande P., Prasad A., Anwar M., Patra. D.D., 2007. Influence of Integrated supply of Vermicompost and Zinc-enriched Compost with two grade levels of iron and zinc on the productivity of Geranium. *Commun Soil Sci Plant Anal.* 38, 2581-2599.
49. Darzi M.T., Ghalavand A., Sefidkon F., Rejali, F., 2008. Effects of vermicompost and biophosphate fertilizer on quantity and quality of essential oil of fennel. *Iran J Med Aroma Plants.* 4(24), 396-413.
50. Fallahi J., Koocheki A., Moghaddam P., 2009. Study the effects of biofertilizers on quantitative and qualitative yield of chamomile. *Iran J Agron.* 7(1), 127-135.
51. Sanches Govin E., Rodrigues Gonzales H., Carballo Guerra C., Milanés Figueredo, M., 2005. Influencia de los abonos organicos y biofertilizantes en la calidad de las especies medicinales *Calendula officinalis* L y *Matricaria recutita* L. *Revista Cubana Plant Med.* 10(1), 1-8.
52. Senthilkumar S., Sriramachandrasekharan M.V., Haripriya K., 2004. Effect of vermicompost and fertilizer on the growth and yield of rose. *J Interacademia.* 8, 207-210.
53. Haj Seyed Hadi M.R., Darzi M.T., Riazi G.H., Ghandehari Z., 2011. Effects of vermicompost and amino acids on the flower yield and essential oil production from *Matricaria chamomile* L. *J Med Plants Res.* 5(23), 5611-5617.